

Amendments to the Claims

1. (Withdrawn) A veterinary composition, useful for providing a rapid onset and long lasting analgesia and sedation in an animal, comprising a pharmaceutically effective amount of a guanidine derivative.
2. (Withdrawn) The composition of claim 1, wherein the guanidine derivative is selected from the group consisting of guanabenz, guanabenz acetate, guanoxabenz, clonidine, guanaciline, guanadrel, guanazodine, guanethidine, guanfacine and guanochlor, guanoxan and chlomidine.
3. (Withdrawn) The composition of claim 1, wherein the guanidine derivative is guanabenz, guanabenz acetate or pharmaceutically acceptable derivatives thereof.
4. (Withdrawn) The composition of claim 1, further comprising a pharmaceutically acceptable carrier.
5. (Withdrawn) The composition of claim 1, wherein the composition is adapted for oral administration.
6. (Withdrawn) The composition of claim 1, wherein the composition is adapted for intravenous administration.
7. (Withdrawn) The composition of claim 1, wherein the composition is adapted for intramuscular administration.
8. (Withdrawn) The composition of claim 1, wherein the animal is selected from the group consisting of equine, canine, feline, bovine, caprine, porcine and ovine.
9. (Withdrawn) The composition of claim 1, wherein the animal is an equine.
10. (Withdrawn) The composition of claim 1, wherein the animal is a standing animal.

11. (Withdrawn) The composition of claim 1, wherein the analgesia and sedation are rapidly reversible.
12. (Withdrawn) The composition of claim 11, wherein the analgesia and sedation are reversed via administration of a pharmaceutically effective amount of an α adrenergic antagonist.
13. (Withdrawn) The composition of claim 12 wherein the α adrenergic antagonist is selected from the group consisting of yohimbine, rauwolscine, idazoxan and atepamezole.
14. (Withdrawn) The composition of claim 1, wherein the pharmaceutically effective amount is between about 0.05 mg/kg and about 0.50 mg/kg.
15. (Withdrawn) The composition of claim 14, wherein the pharmaceutically effective amount is about 0.25 mg/kg.
16. (Withdrawn) The composition of claim 1, wherein the guanidine derivative is guanabenz acetate or a pharmaceutically acceptable derivative thereof and the pharmaceutically effective amount is between about 0.05 mg/kg and about 0.50 mg/kg.
17. (Withdrawn) The composition of claim 1, wherein the guanidine derivative is guanabenz acetate or a pharmaceutically acceptable derivative thereof and the pharmaceutically effective amount is about 0.25 mg/kg.
18. (Withdrawn) The composition of claim 1, wherein the guanidine derivative is an α adrenergic agonist.
19. (Withdrawn) The composition of claim 1 in a unit dosage form.
- 20-64 (Canceled)
65. (New) A method of inducing rapid onset and long lasting sedation and analgesia in an animal, comprising administering to the animal a pharmaceutically effective amount of a composition consisting essentially of a guanidine derivative selected from the group consisting of guanabenz, guanabenz acetate,

guanoxabenz, clonidine, guanaciline, guanadrel, guanazodine, guanethidine, guanfacine, guanochlor, and guanoxan

66. (New) The method of claim 65, wherein the guanadine derivative is guanabenz acetate or pharmaceutically acceptable derivative thereof.

67. (New) The method of claim 65, wherein the administration is oral.

68. (New) The method of claim 65, wherein the administration is intravenous.

69. (New) The method of claim 65, wherein the administration is intramuscular.

70. (New) The method of claim 65, wherein the animal is selected from the group consisting of equine, canine, feline, bovine, caprine, porcine and ovine.

71. (New) The method of claim 65, wherein the animal is an equine.

72. (New) The method of claim 65 wherein the rapid onset sedation and analgesia is induced in a standing animal.

73. (New) The method of claim 65, further comprising the step of selectively reversing or controlling the level of analgesia and sedation in the animal comprising administering a pharmaceutically effective amount of α adrenergic antagonist to the animal.

74. (New) The method of claim 73 wherein the α adrenergic antagonist is selected from the group consisting of yohimbine, rauwolscine, idazoxan and atepamezole.

75. (New) The method of claim 65, wherein the pharmaceutically effective amount of the guanidine derivative is between about 0.05 mg/kg and about 0.50 mg/kg.

76. (New) The method of claim 65, wherein the pharmaceutically effective amount of the guanidine derivative is about 0.25 mg/kg.

77. (New) The method of claim 65, wherein the guanidine derivative is guanabenz acetate or a pharmaceutically acceptable derivative thereof and the pharmaceutically effective amount is between about 0.05 mg/kg and about 0.50 mg/kg.

78. (New) The method of claim 65, wherein the guanidine derivative is guanabenz acetate or a pharmaceutically acceptable derivative thereof and the pharmaceutically effective amount is about 0.25 mg/kg.

79. (New) The method of claim 65, wherein the guanidine derivative is an α adrenergic agonist.

80. (New) A method of inducing rapid onset and long lasting sedation and analgesia in a standing equine animal, comprising administering to the animal a pharmaceutically effective amount of a composition comprised of a guanidine derivative selected from the group consisting of guanabenz, guanabenz acetate, guanoxabenz, clonidine, guanacrine, guanadrel, guanazodine, guanethidine, guanfacine, guanochlor, and guanoxan

81. New) The method of claim 80, wherein the guanidine derivative is guanabenz acetate or pharmaceutically acceptable derivative thereof.

82. (New) The method of claim 80, wherein the administration is oral.

83. (New) The method of claim 80, wherein the administration is intravenous.

84. (New) The method of claim 80, wherein the administration is intramuscular.

85. (New) The method of claim 80, further comprising the step of selectively reversing or controlling the level of analgesia

and sedation in the animal comprising administering a pharmaceutically effective amount of α adrenergic antagonist to the animal.

86. (New) The method of claim 85, wherein the α adrenergic antagonist is selected from the group consisting of yohimbine, rauwolscine, idazoxan and atepamezole.

87. (New) The method of claim 80, wherein the pharmaceutically effective amount of the guanidine derivative is between about 0.05 mg/kg and about 0.50 mg/kg.

88. (New) The method of claim 80, wherein the pharmaceutically effective amount of the guanidine derivative is about 0.25 mg/kg.

89. (New) The method of claim 80, wherein the guanidine derivative is guanabenz acetate or a pharmaceutically acceptable derivative thereof and the pharmaceutically effective amount is between about 0.05 mg/kg and about 0.50 mg/kg.

90. (New) The method of claim 80, wherein the guanidine derivative is guanabenz acetate or a pharmaceutically acceptable derivative thereof and the pharmaceutically effective amount is about 0.25 mg/kg.

91. (New) The method of claim 80, wherein the guanidine derivative is an α adrenergic agonist.